

NuFact08 Perspectives 200






10th International Workshop on Neutrino Factories, Super beams, and Beta beams









2008 Premio Rey Jaime I: Investigación Básica

\$9.5 M for NOVA



NuFact99: Questions of Identity

Do neutrinos oscillate?

What are the neutrino masses?


Is neutrino mass a sign of (nontrivial) BSM physics?

Does the evidence require more than 3 v species?

Can we find evidence for a sterile v?

Could v masses be special (Majorana, scale » EW)?

How could *light* sterile V arise?
Are V mixing angles large? maximal?
Do V masses probe large extra dimensions?
Can we detect CP violation in V mixing?



Three-neutrino mixing formalism

3 V masses, m_1 , m_2 , m_3


$$\mathsf{U} = \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{array}\right) \left(\begin{array}{ccc} c_{13} & 0 & s_{13}e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta} & 0 & c_{13} \end{array}\right) \left(\begin{array}{ccc} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{array}\right)$$

$$\cdot \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & e^{i\varphi_2} & 0 \\ 0 & 0 & e^{i\varphi_3} \end{array} \right) \text{ if } \nu = \bar{\nu} \\ \begin{array}{ccc} & \text{I Dirac phase} \\ \text{2 Majorana phases} \end{array}$$


$$30^{\circ} \lesssim \theta_{12} \lesssim 38^{\circ} \text{ solar}$$
 $35^{\circ} \lesssim \theta_{23} \lesssim 55^{\circ} \text{ atm}$ $\theta_{13} \lesssim 10^{\circ}$

CP phase δ unconstrained


Neutrino family patterns (representative)



Neutrino family patterns (uncertainties)




Neutrino family patterns (tri-bimaximal mixing)




 $\theta_{13} = 0$, $\theta_{23} = \text{maximal}$, $\sin^2 \theta_{12} = \frac{1}{3}$


Quark family patterns



Fermion Masses




Fermion Masses Run



Neutrino Masses: oscillations determine $\Delta m_{\rm V}^2$, not $m_{\rm V}$


$$m_2^2 - m_1^2 \approx 7.9 \times 10^{-5} \text{ eV}^2; |m_3^2 - m_1^2| \approx 2.5 \times 10^{-3} \text{ eV}^2$$




Absolute scale not known, but $m_V \lesssim 10^{-5} m_e$




Relic V contributions to mass density of Universe



Pascoli, Pastor



Interactions: $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ gauge symmetries



Why does the muon weigh? gauge symmetry allows

$$\zeta_e \left[(\overline{e_L} \Phi) e_R + \overline{e_R} (\Phi^{\dagger} e_L) \right] \leadsto m_e = \zeta_e v / \sqrt{2}$$

after spontaneous symmetry breaking

What does the muon weigh?

Ge: picked to give right mass, not predicted

fermion mass implies physics beyond the standard model

How could neutrino mass arise?

Add N_R : SU(2)_L singlet with Y = 0 — sterile couple V_L , N_R in Dirac mass term, $C_V \lesssim 10^{-11}$

V has no charge or color, so $V \equiv \overline{V}$ is possible (Majorana)

Dirac & Majorana: explain m_V ?

Neutrino mass as physics beyond standard model: may connect with ultrahigh scales (seesaw) but might also implicate TeV scale

The Importance of the I-TeV Scale

EW theory does not predict Higgs-boson mass Thought experiment: conditional upper bound

 $W_L^+W_L^-, Z_L^0Z_L^0, HH, HZ_L^0$ satisfy $\emph{s}\text{-wave}$ unitarity,

provided
$$M_H \leq \left(8\pi\sqrt{2}/3G_F\right)^{1/2} = 1 \text{ TeV}$$

- If bound is respected, perturbation theory is everywhere reliable
- If not, weak interactions among W^{\pm} , Z, H become strong on I-TeV scale

New phenomena are to be found around I TeV





If electroweak symmetry were not hidden ...

- Massless quarks and leptons
- QCD confines quarks into color-singlet hadrons
- Nucleon mass little changed
- QCD breaks EW symmetry, gives tiny W, Z masses; weak-isospin force doesn't confine
- p outweighs n: rapid β-decay
 - \Rightarrow lightest nucleus is $n \dots$ no hydrogen atom
- ullet Some light elements from BBN, but ∞ Bohr radius
- No atoms means no chemistry, no stable composite structures like liquids, solids, ...

... character of the physical world would be profoundly changed

Parameters of the Standard Model

```
coupling parameters \alpha_s, \alpha_{\rm em}, \sin^2 \theta_W
      parameters of the Higgs potential
      vacuum phase (QCD)
                                Flavor physics may be where we see, or diagnose,
6
      quark masses
3
      quark mixing angles
                                the break in the SM.
1
3
3
1
      CP-violating phase
      charged-lepton masses
      neutrino masses
      leptonic mixing angles
      Ieptonic CP-violating phase (+ Majorana ...)
      arbitrary parameters
```

What is the nature of the mysterious new force that hides electroweak symmetry?

- *A force of a new character, based on interactions of an elementary scalar
- *A new gauge force, perhaps acting on undiscovered constituents
- *A residual force that emerges from strong dynamics among electroweak gauge bosons
- *An echo of extra spacetime dimensions


Essential step toward understanding the new force that shapes our world:

Find the Higgs boson and explore its properties.

- * Is it there? How many?
- * Verify quantum numbers (spin, parity, ...)
- * Does H generate mass for gauge bosons and for fermions?
- * How does H interact with itself?

New Physics on the Fermi Scale?

If dark matter interacts weakly ...




... its likely mass is 0.1 to 1 TeV: Fermi scale

How to separate EW, higher scales?

Does M_H < I TeV make sense?

The peril of quantum corrections — hierarchy problem



How to separate EW, higher scales?

Traditional: change electroweak theory to understand why M_H , electroweak scale $\ll M_{Planck}$

To resolve hierarchy problem: extend standard model on the I-TeV scale ...

 $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$

composite Higgs boson technicolor / topcolor supersymmetry

• • •


Ask instead why gravity is so weak, why $M_{Planck} \gg$ electroweak scale

extra dimensions

Look forward to immensely productive conversations among LHC discoveries, neutrino advances, other high-sensitivity accelerator experiments, and astro/cosmo/particle observations

Many examples of possible connections presented at NuFact08

Lifting the electroweak veil around I TeV should help us to see the problem of identity (flavor) and the challenges of other scales more clearly



New physics in the standard model?

What phenomena are implied by the standard model, but too subtle to have attracted our notice?

Known example: sphaleron for B-violation

Recent case: $Z\gamma\omega$ anomaly-mediated $v\gamma$ interaction in presence of baryons (Harvey, Hill, & Hill)

Added motivation to measure & understand V cross sections, hadroproduction at low energies HARP, MIPP, SciBooNE, MinerVa, MiniBooNE, T2K, ... Catanesi, Harris, Nakaya, Petti, ...



Enormously impressive progress reported here

Coherence of experimental results Breadth of scientific opportunities Creativity of machine designers

Much incentive to stretch our minds, think "blue-sky" thoughts about machines, detectors, baselines, strategies ...

and even to ask how little we could require

Super Beams



♦ J-PARC facility in Japan:

- Phase I for T2K (2009): 0.75 MW 30 GeV, 10×10^{20} pot/y;
- Phase II upgrade (2015): $4MW \ 50 \ GeV, \ 50 \times 10^{20} \ pot/y$




◆ Project-X at Fermilab:

- NuMI for NO ν A: 400kW, 700kW MI 120 GeV, 6×10^{20} pot/y;
- New GeV linac + recycler + MI (2016): 2.3 MW 120 GeV, 30×10^{20} pot/y.


(see talk by D. Harris & A. Jansson)

Neutrino Factory



- **♦** Megawatt proton source of few GeV for pion production
- ◆ Neutrinos produced from decay of 25-50 GeV stored muons
 - \implies Useful decays achievable $3 \div 5 \times 10^{20}$ per year

β Beams



- ♦ Megawatt proton driver of few GeV to produce radioactive ions (ISOL target)
- igspace Neutrinos originated from β decay of accelerated ions (EURISOL)
 - \Longrightarrow Rates of $2.9 \times 10^{18} \bar{\nu}$ (6 He) and $1.1 \times 10^{18} \nu$ (18 Ne) per year at $\gamma = 100$


+ 8Li, 8B, EC monochromatic beams

Wildner

From existing detectors to a new generation

Magnetized iron
Water Cherenkov
Magnetized emulsion
Totally active scintillator
Liquid argon

• •



Practical considerations will intervene

Cavern size, structural integrity, excavation time

Detector cost, fabrication time

Difficulty, cost of steep dip angles

What compromises will get us to the essential physics results in the shortest time?

Realizing Neutrino Factory or B Beam

When will needed demonstrations be in hand? When will experimental specification be needed?

When might we hope to ask for funding? How long to operations?

Bagnères-de-Bigorre, 1953 the "last" cosmic-ray conference the coming of the BNL Cosmotron

C. F. Powell: "Gentlemen, we have been invaded!

The accelerators are here."

Louis Leprince-Ringuet: « Mais nous devons aller vite, nous devons courir sans ralentir notre cadence : nous sommes poursuivis ... nous sommes poursuivis par les machines! »

NuFact08: Updated Questions of Identity

What are subdominant neutrino transitions?
What is the neutrino hierarchy, absolute mass scale?
Is neutrino mass a sign of nontrivial BSM physics?
Bottom line on LSND & MiniBooNE?
Can we find evidence for a sterile v?
Dirac or Majorana (L-number), scale >> EW, GUTs?

Connection between V mass, LFV, what else? How could light sterile V arise? θ_{23} : is V_3 mostly V_μ or V_τ ? How small is θ_{13} ? Do V masses probe large extra dimensions? Can we detect CP violation in V mixing? Does leptogenesis explain matter excess? How do neutrinos shape the universe? V decay, EDM, MDM, discrete symmetry violation?

Thanks to all!